



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Mortuary report.—Congenital debility, 1; unknown, 1; tuberculosis, pulmonary, 2; pernicious fever, 1 (a child of $2\frac{1}{2}$ years); cardiac insufficiency, 1; pneumonia, 1; sudden death, 1; cirrhosis of the liver, 1; total, 9. Mortality rate, 23.80. Sanitary situation: The mortality rate during the past week was one of the lowest.

Reports from Veracruz—*Yellow-fever infection probably imported—Stegomyia increasing in numbers and area of diffusion.*

Passed Assistant Surgeon Wilson reports, July 12, as follows:

One case of yellow fever was reported in Veracruz on the 11th and 2 cases on the 12th instant. The infection of one, if not more, of these cases originated in Veracruz. Owing to the large number of nonimmunes present in the city (about 12,000, more than one-third of the population), it is to be feared that many more cases will develop.

Week ended July 15, 1905: Vessels inspected and bills of health issued, 9; vessels fumigated with sulphur to kill mosquitoes, 3; crew inspected, 325; passengers inspected, 96.

Three cases of yellow fever were reported during the week, but no deaths. These cases were infected in Veracruz, the infection coming originally, in all probability, from Tierra Blanca, where cases have been almost continuously under treatment for several weeks past and where there are at present two cases. *Stegomyia* are increasing in numbers and appear to be more widely diffused than they were a few weeks ago. Rains are not frequent.

Yellow-fever cases.

VERACRUZ, MEXICO, July 24, 1905.

WYMAN, Washington:

Three cases yellow fever, 23d and 24th.

WILSON.

VERACRUZ, MEXICO, July 26, 1905.

WYMAN, Washington:

One case yellow fever, July 25.

WILSON.

NICARAGUA.

Report from Bluefields, fruit port—*Case of elevated temperature noted on steamship Bluefields.*

Acting Assistant Surgeon Layton reports as follows: Week ended July 11, 1905. Present officially estimated population, 3,500; 4 deaths; prevailing disease, malarial fever; general sanitary condition of this port and the surrounding country during the week, good. Captain of steamship *Bluefields* when ship left had temperature of $102\frac{2}{5}^{\circ}$. During week in port captain's temperature ranged between 100° and 103° daily.

Allowed shipping of one dead body July 6.

Deaths, 1 from epilepsy, 3 accidental, from drowning.